

ALERTS

Timely Information Directly to Your Inbox

DynaMed Plus®

There's no need to search for the latest development; let *DynaMed Plus*® notify you. Sign up to get alerts about the latest clinical and practice-changing news and information.

Sign up for alerts of recent updates by category or, more specifically, by *Practice Changing Updates*. **Click Follow**, set the frequency, and receive an email notification when recent updates are made.

EBSCOHealth Calculators Sign Up For Remote Access Feedback Help About

DynaMed Plus Search DynaMed Plus Search

Recent Updates

All Categories Practice Changing Updates Only Follow

08/24/2015 06:52:00 AM (ET)
probiotics may reduce risk of eczema or atopic eczema in infants (J Allergy Clin Immunol 2015 Jun 1 early online)
Topic: Probiotics, Atopic dermatitis

08/24/2015 06:49:00 AM (ET)
platelet-rich plasma injection may slightly reduce pain at 3 months compared to corticosteroid injection in patients with plantar fasciitis
Topic: Plantar fasciitis

08/24/2015 06:47:00 AM (ET)
small group balance and exercise training might reduce risk of injurious falls in community-dwelling women aged 75-85 years w
Topic: Falls in the elderly

08/24/2015 06:45:00 AM (ET)
ibuprofen 400 mg associated with increased pain-free rate at 2 hours in adults with frequent episodic tension-type headache (C
Topic: Tension-type headache

08/24/2015 06:44:00 AM (ET)
oral cytology may have high sensitivity and specificity for diagnosing oral squamous cell carcinoma or potentially malignant dis
Database Syst Rev 2015 May 29
Topic: Head and neck cancer

08/24/2015 06:43:00 AM (ET)
addition of larazocin 0.5 mg to gluten-free diet may improve symptoms in patients with celiac disease (Gastroenterology 2015
Topic: Celiac disease

EBSCOHealth Calculators Sign Up For Remote Access Feedback Help About

DynaMed Plus Search DynaMed Plus Search

Recent Updates

All Categories Practice Changing Updates Only

08/24/2015 06:52:00 AM (ET)
probiotics may reduce risk of eczema or atopic eczema in infants (J
Topic: Probiotics, Atopic dermatitis

08/24/2015 06:49:00 AM (ET)
platelet-rich plasma injection may slightly reduce pain at 3 months o
Topic: Plantar fasciitis

08/24/2015 06:47:00 AM (ET)
small group balance and exercise training might reduce risk of injur
Topic: Falls in the elderly

08/24/2015 06:45:00 AM (ET)
ibuprofen 400 mg associated with increased pain-free rate at 2 hours in adults with frequent episodic tension-type headache (Cochrane Database Syst Rev 2015 Jul 31)
Topic: Tension-type headache

Follow

All Categories

Your E-mail:

Email Updates:

Once a day
Once a week
Once a month

Submit

Sign up for alerts linked to any topic of interest. **Click Follow**, set the frequency, and receive an email notification when there is a notable change in clinical content.

EBSCOHealth Calculators Sign Up For Remote Access Feedback Help About

DynaMed Plus Sepsis treatment in adults Search

Sepsis Updates

Related Summaries

Overview

- Initial Supportive Care
- Antibiotics and Source Control
- Corticosteroids
- Mechanical Ventilation
- Renal Replacement Therapy
- Glucose Control
- Other Supportive Care
- Intravenous Immunoglobulin (IVIg)
- Therapies with Limited Evidence or Therapies No Longer Recommended
- Quality Improvement
- Guidelines and Resources
- Patient Information
- ICD-9/ICD-10 Codes
- References

Follow

Sepsis treatment in adults

Your E-mail:

Email Updates:

Once a day
Once a week
Once a month

Submit

From: epalerts@epnet.com [mailto:epalerts@epnet.com]
Sent: Tuesday, April 12, 2016 2:40 AM
To: Jane Dough <JDough@EBSCO.COM >
Subject: DynaMed Plus: Following Sepsis treatment in adults

DynaMed Plus

Following: **Sepsis treatment in adults**
Daily Update Apr 12, 2016

- Apr 11, 2016 8:06 AM**
Temporary fluid response induced by passive leg raising may help identify critically ill patients responsive to fluid therapy (Crit Care Med 2016 Jan 6 early online)
Topics: [Sepsis treatment in adults](#)
- Apr 11, 2016 7:07 AM**
Prolonged low-dose corticosteroids may reduce 28-day mortality in patients with sepsis but corticosteroids given at any dose associated with increase in hyperglycemia and hypenatremia (Cochrane Database Syst Rev 2015 Dec 3)
Topics: [Sepsis treatment in adults](#), [Critical illness-related corticosteroid insufficiency](#)

To view this update, you may need login credentials. If you have them, you can access directly via the link(s) above. If you need login credentials, please contact your librarian or DynaMed Plus administrator.